Author Index

Ackermand, D., Herd, R. K., Reinhardt, M. & Windley, B. F. Sapphirine parageneses from the Caraiba complex, Bahia, Brazil: the influence of Fe^{2+}–Fe^{3+} distribution on the stability of sapphirine in natural assemblages, 323.

Aguirre, L. & Atherton, M. P. Low-grade metamorphism and geotectonic setting of the Macuchi Formation, Western Cordillera of Ecuador, 473.

Atherton, M. P., *see* Aguirre, L.

Baker, A. J. Models for the tectothermal evolution of the eastern Dalradian of Scotland, 101.

Baker, J., Powell, R., Sandiford, M. & Muhling, J. Corona textures between kyanite, garnet and gedrite in gneisses from Errabiddy, Western Australia, 357.

Bargossi, G. M., *see* Morten, L.

Baroz, F., Bebien, J. & Ikenne, M. An example of high-pressure metamorphic rocks from an island-arc: the Paikon Series (Innermost Hellendides, Greece), 509.

Bebien, J., *see* Baroz, F.

Black, L. P., Harley, S. L., Sun, S. S. & McCulloch, M. T. The Rayner Complex of East Antarctica; complex isotopic systematics within a Proterozoic mobile belt, 1.

Black, P. M., *see* Ghent, E. D.

Brothers, R. N., *see* Ghent, E. D.

Bucher-Nurminen, K., *see* Klaper, E. M.

Burg, J.-P., *see* Clarke, G. L.

Burg, J.-P., *see* Guiraud, M.

Clarke, G. L., Guiraud, M., Powell, R. & Burg, J.-P. Metamorphism in the Olary Block, South Australia: compression with cooling in a Proterozoic fold belt, 291.

Craw, D., *see* Jamieson, R. A.

Daly, J. S., *see* Sanders, I. S.

D'Arcy, W. F., *see* Vernon, R. H.

Davies, G. R., *see* Sanders, I. S.

Diaz de Federico, A. *see* Morten, L.

Emmett, T. F. A reconnaissance study of the distribution of Ba, Nb, Y, and Zr in some Jotun kindred gneisses from Central Jotunheimen, southern Norway, 41.

Ferrati, N., *see* Pognante, U.

Flood, R. H., *see* Vernon, R. H.

Ghent, E. D., Stout, M. Z., Black, P. M. & Brothers, J. Chloritoid-bearing rocks associated with blueschists and eclogites, northern New Caledonia, 239.

Guiraud, M., Burg, J.-P. & Powell, R. Evidence for a Variscan suture zone in the Vendée, France: a petrological study of blueschist facies rocks from Boi de Cené, 225.

Guiraud, M., *see* Clarke, G. L.

Harley, S. L. A pyroxene-bearing meta-ironstone and other pyroxene-granulites from Tonagh Island, Enderby Land, Antarctica: further evidence for very high temperature (>980°C) Archaean regional metamorphism in the Napier Complex, 341.

Harley, S. L. *see* Black, L. P.

Hammond, R. L. The influence of deformation partitioning on dissolution and solution transfer in low-grade tectonic mélange, 195.

Hensen, B. J. *P–T* grids for silica-understaurated granulites in the system MAS (n+4) and FMAS (n+3)—tools for the derivation of *P–T* paths of metamorphism, 255.

Hensen, B. J., *see* Warren, R. G.

Herd, R. K., *see* Ackermand, D.

Ikenne, M., *see* Baroz, F.

Itaya, T., *see* Okuyama-Kusunose, Y.

Jamieson, R. A. & Craw, D. Sphalerite geobarometry in metamorphic terranes: an appraisal with implications for metamorphic pressure in the Otago schist, 87.

Jamieson, R. A. & Vernon, R. H. Timing of porphyroblast growth in the Fleur de Lys Supergroup, Newfoundland, 273.

Keskinen, M. & Liou, J. G. Stability relations of Mn–Fe–Al piemontite, 495.

Kienast, J. R., *see* Vernié, P.

Klaper, E. M. & Bucher-Nurminen, K. Alpine metamorphism of pelitic schists in the Nufenen Pass area, Lepontone Alps, 175.

Leclair, A. D., *see* Thompson, P. H.

Liou, J. G., *see* Keskinen, M.

Liou, J. G. & Maruyama, S. Parageneses and compositions of amphiboles from Franciscan jadeite–glaucophane type facies metabasites at Cazadero, California, 371.

Liou, J. G., *see* Maruyama, S.

Martinez Martinez, J. M., *see* Morten, L.

Maruyama, S. & Liou, J. G. Clinopyroxene—a mineral telescoped through the processes of blueschist facies metamorphism, 529.

Maruyama, S., *see* Liou, J. G.

McCulloch, M. T. *see* Black, L. P.

Mevel, C., *see* Vernié, P.

Morten, L., Bargossi, G. M., Martinez Martinez, J. M., Puga, E. & Diaz de Federico, A. Metagabbro and associated eclogites in the Lubrin area, Nevado-Filabride Complex, Spain, 155.

Muhling, J., *see* Baker, J.

Okay, A. I. The oxygen fugacity stability of deerite: an alternative view, 553.

Okuyama-Kusunose, Y. & Itaya, T. Metamorphism of carbonaceous material in the Tono contact

aureole, Kitakami Mountains, Japan, 121.
Phillips, G. N. Metamorphism of the Witwatersrand gold fields: conditions during peak metamorphism, 307.
Pognante, U., Talarico, F., Rastelli, N. & Ferrati, N. High pressure metamorphism in the nappes of the Valle dell'Orco traverse (Western Alps collisional belt), 397.
Powell, R., *see* Baker, J.
Powell, R., *see* Clarke, G. L.
Powell, R., *see* Guiraud, M.
Prior, D. J. Syntectonic porphyroblast growth in phyllites: textures and processes, 27.
Puga, E., *see* Morten, L.
Rastelli, N., *see* Pognante, U.
Reinhardt, J. Cordierite–anthophyllite rocks from north-west Queensland, Australia: metamorphosed magnesian pelites, 451.
Reinhardt, M., *see* Ackermand, D.
Ryburn, R. J., *see* Warren, R. G.
Sanders, I. S., Daly, J. S. & Davies, G. R. Late Proterozoic high-pressure granulite facies metamorphism in the north-east Ox inlier, north-west Ireland, 69.
Sandiford, M., *see* Baker, J.
Snow, E. & Yund, R. A. The effect of ductile deformation on the kinetics and mechanisms of the aragonite—calcite transformation, 141.
Stout, M. Z., *see* Ghent, E. D.
Sun, S. S. *see* Black, L. P.
Talarico, F., *see* Pognante, U.
Thompson, P. H. & Leclair, A. D. Chloritoid–hornblende assemblages in quartz–muscovite pelitic rocks of the Central Metasedimentary Belt, Grenville Province, Canada, 437.
Vernié, P., Kienast, J. R. & Mevel, C. Deerite in highly oxidizing conditions: a reply, 557.
Vernon, R. H. Growth and concentrations of fibrous sillimanite related to hetergenous deformation in K-feldspar–sillimanite metapelites, 51.
Vernon, R. H., Flood, R. H. & D'Arcy, W. F. Sillimanite and andalusite produced by base–cation leaching and contact metamorphism of felsic igneous rocks, 439.
Vernon, R. H., *see* Jamieson, R. A.
Warren, R. G., Hensen, B. J. & Ryburn, R. J. Wollastonite and scapolite in Precambrian calc-silicate granulites from Australia and Antarctica, 213.
Windley, B. F., *see* Ackermand, D.
Yund, R. A., *see* Snow, E.

Subject Index

actinolite, 371
Alpine metamorphism, 175
aluminosilicate polymorphs, 291
andalusite, 439
Andes, Ecuador, 473
anisotropic domain textures, 121
Antarctica, 1, 341
aragonite–calcite transformation kinetics, 141
Archaean, 341

base–cation leaching, 51, 439
Betic Cordilleras, Spain, 155
blueschist, 141, 225, 529
blueschist facies, 509

calc-silicate rocks, 213
Canadian shield, 415
coaly material, 121
chloritic alteration, 451
chloritoid, 225, 239, 307
chloritoid–hornblende assemblages, 415
clinopyroxene, 529
collisional belt, 397
contact metamorphism, 121, 439
cordierite, 357
cordierite–anthophyllite rocks, 451
cordierite stability, 255
corona textures, 357
crossite, 371
crustal imbrication, 69
crystal fractionation, 41

deformation-enhanced reactions, 141
deformation mechanism, 141
deformation partitioning, 27, 195

eastern Dalradian, 101
eclogite, 69
epidote minerals, 495
evaporitic clays and shales, 451
experimental petrology, 495
felsic hornfelses, 439
fibrous sillimanite, 51
Fleur de Lys Super Group, Newfoundland, 273
fluid circulation, 155
fluid flow, 195
fluid-phase composition, 213
foliation, 51

gabbro–eclogite transformation, 155
garnet, 273, 357
garnet–chlorite thermometer, 239
gedrite, 357
geobarometry, 87, 371
geochemistry, 69
geochronology, 1
geotectonic setting, 473
geothermobarometry, 69, 175
glaucophane, 225, 371
gold, 307
granulite facies, 323
granulites, 41, 213, 255
greenschist, 307
greenschist facies, 509
Grenville Province, 415

Hellenides, Greece, 509
Helvetic–Penninic contact zone, 175
heterogeneous deformation, 51
high-pressure granulite facies, 69
high-pressure metamorphism, 239, 397
high-temperature graphitization, 121

inclusion trails, 273
Ireland, 69
island arc, 473
isobaric cooling, 69
isograds, 175
isotopic resetting, 1

Jotunheimen, Norway, 41

kyanite zone, 101

lawsonite, 509
Lepontine Alps, 175
LIL and HFS trace elements, 41
low-grade metamorphism, 473

marginal basin, 473
metabasite, 529
metamorphism, 341
Mount Isa Block, Australia, 451

Napier Complex, 341
New Caledonia, 239
north-east Ox inlier, 69
nucleation, 27

Olary Block, Willyama Complex, 291
Otago schist, 87
oxygen fugacity, 495

peak *T*, 255
pelite metamorphism, 175
perturbed geotherm, 291
petrogenetic grids, 323, 415
phyllite, 27
piemontite, 495
pigeonite, 341
porphyroblast, 273
porphyroblast growth, 27
porphyroblast rotation, 27

P–T grids, 255
P–T path, 255, 291, 495
P–T–t path, 69
Proterozoic, 1, 69, 451
pyrophyllite, 307

Rayner Compex, 1
reaction kinetics, 397
riebeckite, 371

S and C planes, 195
sapphirine, 323, 357
scapolite, 213
sillimanite, 439
sillimanite zone, 101
Sm–Nd, 69
solution transfer, 195
sphalerite, 87
sulphide metamorphism, 87
sulphides, 439

tectonic mélange, 195
tectonic thickening, 101
textures, 27
thrust tectonics, 397
timing of growth, 273

Vendée, France, 225
very high temperature, 341

Ward Creek, Cazadero, 371, 529
Western Alps, Italy, 397
whole rock geochemistry, 41
winchite, 371
Witwatersrand, 307
wollastonite, 213